

Scotland's Rural College

## Hydrazone comprising compounds as promising anti-infective agents

Sharma, P. C.; Sharma, D.; Sharma, A.; Saini, N.; Goyal, R.; Ola, M.; Chawla, R.; Thakur, V. K.

*Published in:*  
Materials Today Chemistry

*DOI:*  
[10.1016/j.mtchem.2020.100349](https://doi.org/10.1016/j.mtchem.2020.100349)

Print publication: 01/12/2020

*Document Version*  
Peer reviewed version

[Link to publication](#)

*Citation for pulished version (APA):*  
Sharma, P. C., Sharma, D., Sharma, A., Saini, N., Goyal, R., Ola, M., Chawla, R., & Thakur, V. K. (2020). Hydrazone comprising compounds as promising anti-infective agents: chemistry and structure-property relationship. *Materials Today Chemistry*, 18, [100349]. <https://doi.org/10.1016/j.mtchem.2020.100349>

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.


- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

### Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

**Highlights:**

- 1) This review highlights the potential of hydrazone derivatives as the development of newer pharmacologically active anti-infective agents.
- 2) SAR study of hydrazone based analogues and concise description of nanocarriers as efficient anti-infective agents.
- 3) A wide range of analogues has potent anti-infective activity against various bacterial, fungal and viral strains.
- 4) A brief discussion on patents filed/granted on hydrazone derivatives as anti-infective agents.



## **Hydrazone Comprising Compounds as Promising Anti-infective Agents: Chemistry and Structure-Property Relationship**

**Prabodh Chander Sharma<sup>1\*</sup>, Diksha Sharma<sup>2</sup>, Archana Sharma<sup>2</sup>, Neha Saini<sup>2</sup>, Rajat Goyal<sup>2</sup>, Monika Ola<sup>3</sup>, Rakesh Chawla<sup>4</sup>, Vijay Kumar Thakur<sup>5,6\*</sup>**

<sup>1</sup>Department of Pharmaceutical Chemistry, Delhi Pharmaceutical and Research University, New Delhi-110017

<sup>2</sup>Institute of Pharmaceutical Sciences, Kurukshetra University, Kurukshetra-136119, India,

<sup>3</sup>Department of Pharmaceutics, R.C. Patel Institute of Pharmaceutical Education and Research, Shirpur-425405, India

<sup>4</sup>Department of Pharmaceutical Chemistry, University Institute of Pharmaceutical Sciences and Research, Baba Farid University of Health Sciences, Faridkot-151203, India

<sup>5</sup>Biorefining and Advanced Materials Research Center, Scotland's Rural College (SRUC), Kings Buildings, Edinburgh, EH9 3JG, UK

<sup>6</sup>Department of Mechanical Engineering, School of Engineering, Shiv Nadar University, Uttar Pradesh, 201314, India

### **Abstract**

Despite the adverse effects of microbial hazards on public health, major pharmaceutical firms have left the field of anti-infective development and a dramatic reduction in the number of researcher's intricated in the quest for new specific anti-infective leads. In the non-existence of an efficient forum for antibiotics development and over usage in human beings and animals, bacteria have demoralized this potential by gradually establishing the resistance toward most of the antibiotics used. Thus, the production of novel and successful anti-infective drugs is urgently required to combat this resistance. Hydrazones and hydrazides have mounted as a key-skeleton for the development of active drugs, due to their important biological and pharmacological profiles. Hydrazones are being manufactured as medicines by various investigators to fight against the ailments with maximal effects and minimal toxicity. This paper focuses on the outline of the literature results of recent years, incorporating the work on the anti-infective profile of

hydrazonanalogue. This review may also act as an excellent basis for the development of new derivatives of hydrazone as potential anti-infective mediators.

**Keywords:** Hydrazones, anti-infective, antiviral, antitubercular, biological activity, structure-activity relationship.

**Highlights:**

- 1) This review highlights the potential of hydrazone derivatives as the development of newer pharmacologically active anti-infective agents.
- 2) SAR study of hydrazone based analogues and concise description of nanocarriers as efficient anti-infective agents.
- 3) A wide range of analogues has potent anti-infective activity against various bacterial, fungal and viral strains.
- 4) A brief discussion on patents filed/granted on hydrazone derivatives as anti-infective agents.

## CONTENTS:


1.	Introduction	4-10
2.	Structure-activity relationship study of hydrazones	10-11
3.	Synthesis via conventional and green chemistry	11-13
4.	Literature survey of hydrazones	13-33
4.1	Antimicrobial activity	13-20
4.2	Antiviral Activity	21-25
4.3	Anti-TB Activity	25-33
5.	Recent patents filed/granted on hydrazone derivatives	34-35
6.	Conclusion	35
7.	Future Aspects	36
	Conflict of interest	36
	Abbreviations	36-37
	References	37-49

## 1. INTRODUCTION

Hydrazone ligands have attracted particular attention from researchers due to their well-known chelating capability, structural versatility and diverse range of applications. The metal complexes of hydrazones are including heterocyclic moieties containing nitrogen, oxygen and sulphur as hetero-atoms have been studied extensively to create a probable relationship between the chemical structure and biological activity. Among them, Schiff's base hydrazones bearing nitrogen-containing moiety have attracted considerable attention due to their impressive chemical and physical properties, biological activities and also their analytical applications [1]. They are considered as the major category for the development of new drugs, determination of innovative biologically active molecules against multidrug-resistant microbial infections and can act as multidentate ligands with transition metals which demonstrate many therapeutic applications [2]. Hydrazone characterizes as a resourceful compound having a basic structure  $R^1R^2C=NNR^3R^4$ . Azomethine is a very important group of Schiff's bases, in which  $C=N$  is used as interesting ligand in coordination chemistry [3-5]. Hydrazone-type compounds containing an azomethine constitute an important class of compounds for the production of new drugs. The hydrazone community is well known to play a significant role in anti-infective profile [6]. Hydrazones are related to the ketones and aldehydes. They are formed by substituting the oxygen group of carbonyl compounds with the functional group  $-NNH_2$ . They act as reactants in several important reactions such as hydrazone iodination, Shapiro reaction and Bamford-Stevens reaction to form vinyl compounds. Also, they act as an intermediate in Wolff-Kishner reduction [7]. Hydrazones are mainly synthesized by refluxing the appropriate quantity of substituted hydrazines/hydrazides with ketones and aldehydes in appropriate solvents like tetrahydrofuran, glacial acetic acid, methanol, ethanol, butanol, ethanol-glacial acetic acid etc. Coupling of aryl diazonium salts with active hydrogen compounds can also synthesize hydrazones [8]. Hydrazides and hydrazones are not only intermediates but, in their own right, they are also very effective organic compounds [9]. They contain nitrogen atoms which have nucleophilic and the carbon atom has both nucleophilic and electrophilic effect. The hydrazones endowed with  $\alpha$ -hydrogen are more potent as compared to acidic ketones. Schiff's bases produced by salicylaldehyde and its analogues with primary amines, bearing the  $N_2O$ ,  $NO_2$ ,  $N_2S$  or  $NSO$  donor sets, possess scrupulous biological activities [10-12]. They were commonly used in the identification and quantitative determination of various metallic compounds, analytical chemistry for the detection

and isolation of carbonyl compounds, for preparation of compounds with different structures. However, their greater physiological activity is the most important property of hydrazone derivatives [13]. Nowadays, emerging bacterial resistance is the chief difficulty in the handling of several infections, with the intention of that many hydrazone compounds have been prepared and examined for their anti-infective activity [14-15].

Recently, metal complexes of Schiff's bases have been fascinated the considerable interest due to their versatile pharmacological activities described in **Fig. 1**. such as anti-cancer [16-17], antiviral [18-19], anti-TB [20-21], antimicrobial [22-23], anti-inflammatory [24-25], anticonvulsant [26-27], antiprotozoal [28-29], anti-HIV [30-31], anti-oxidant [32] and analgesic [33-34]. Few marketed drugs bearing hydrazone nucleus related to various biological activities are mentioned in **Fig. 6**.




**Fig.1.** Various pharmacological activities of hydrazone derivatives.

Nanotechnology generally refers to an area of applied science and technology whose general theme is atomic and molecular control of matter [35]. Nanotechnology in medicine, particularly in drug delivery is already widely used and discussed in many medical fields. Not only the drug can be distributed proficiently to the site of infection with the aid of nanomaterials but even the




quantity and intensity of dosage can be regulated by avoiding therapy-related toxicity [36]. Nanoparticles are materials with at least one dimension (1-100 nm) within the range of the nanometer scale or whose basic unit is within this range in three-dimensional space (various applications, types and properties of nanoparticles are shown in **Fig. 2**).




**Fig. 2.** Applications, types and properties of nanoparticles [37,38].

In particular, nanoparticles have a wide range of antibacterial properties towards Gram-positive and Gram-negative bacteria [39]. Because of its high surface area to volume ratio and specific physical and chemical properties, inorganic compounds in nanosize show significant antibacterial activity at low concentrations [40]. The bactericidal activity of these nanocarriers depends greatly on stability, the concentration of the growth medium and size [41]. When immersed in the biological cultural medium, nanocarriers experience various biological interfaces due to the presence of cellular molecules such as proteins, DNA, lipids, polysaccharides and flavonoids etc


[42]. Mechanism of action of nanoparticles based hydrazones as antibacterial agents is depicted in **Fig. 3**. Viruses present a rather significant threat to the medical, pharmaceutical and biotechnological fields as a key factor of sickness and mortality of humans. The extraordinary capacity of viruses to change easily at the new host and transitioning to a different host is really troubling and a big recovery shortage. Nanotechnology provides a platform that modifies or develops the drugs into nanoparticles for the treatment of various infectious diseases. As nanomedicines have smaller size they have ability to interact easily with the viruses or other microbes [43]. Mechanism of action of nanoparticles based hydrazone compounds as antiviral agents is depicted in **Fig. 4**. A millions of people are affected by tuberculosis and it is the second leading cause of death from infectious diseases after AIDS. Nanotechnology offers a new and effective strategy to develop antimycobacterial formulations [44]. The mechanism of action of nanoparticles based hydrazone compounds as anti-tubercular agents have shown in **Fig. 5**.




**Fig. 3.** Mechanism of action of nanoparticles based hydrazones compounds as antibacterial agents [45,46].



**Fig. 4.** Mechanism of action of nanoparticles based hydrazones compounds as antiviral agents [47,48].



**Fig. 5.** Mechanism of action of nanoparticles based hydrazone compounds as anti-tubercular agents [49,50].




**Fig. 6.** Commonly prescribed marketed drugs bearing hydrazone nucleus [51-54].

## 2. STRUCTURE ACTIVITY RELATIONSHIP STUDY OF HYDRAZONES

The structure-activity relationship is the relationship among the 3D-structure or chemical structure of a molecule and its pharmacological activity. The SAR analysis allows determining the chemical groups accountable for triggering a biological target influence in the organism. Medicinal chemists utilize chemical synthesis methodologies to introduce new chemical groups into medicinal compounds and check its modification for their pharmacological impacts [55].

Hydrazone nucleus comprises N-NH<sub>2</sub> functional group, the substitution of R<sub>1</sub> with phenyl ring containing bromine, 3-methoxy-4-hydroxy groups showed superior antimicrobial activity in most of the analogues [56]. In some derivatives, if R<sub>1</sub> is substituted with 5-nitrofur-2-yl compounds showed better antibacterial activity [57]. Substitution of R<sub>1</sub> with electron-accepting groups like

F, Cl, Br and NO<sub>2</sub> groups displayed strong antifungal activity while analogues with heterocyclic moieties like thiophene and indole demonstrated good antimicrobial activity [58]. The most potent antitubercular activity was observed when R<sub>1</sub> is substituted with chlorine, bromine and nitro groups [59]. It was also found that if R<sub>2</sub> is substituted with benzothiazole it exhibited higher antimicrobial as well as iron-chelating activity [60]. Another substitution of R<sub>2</sub> with thiazole-4-(p-halophenyl) increases antibacterial activity [61]. When R<sub>1</sub> was substituted with hydroxyl and methoxy group compound showed superior antimicrobial activity [62]. Electron donating substituents on phenyl ring at R<sub>2</sub> displayed moderate resistance toward microbes [63]. The SAR of hydrazones as anti-infective agents is detailed in **Fig. 7**.




**Fig. 7.** SAR of the anti-infective potential of hydrazone derivatives.

### 3. SYNTHESIS VIA CONVENTIONAL AND GREEN CHEMISTRY

Over the decades the conventional chemical synthesis utilizes the hazardous and toxic chemicals and solvents in the manufacturing of by-products. This is an intentional implementation of chemical reactions to produce one or more products. To overcome all these unfavourable issues green chemistry came into consideration with an innovative approach for the synthetic chemist to

obtain the product with maximum yield as eco-friendly. The green chemistry approaches were observed as the best economic way to obtain a maximum number of synthetic molecules in desirable concentration. This chemical synthesis explores our potential challenges in dealing with chemical processes and products by introducing new reactions that can optimize the desired products and reduce by-products, developing new synthetic schemes and instruments that can improve chemical synthesis and searching green solvents that are eco-friendly. Microwave-assisted synthesis (MAOS) is considered as a synthetic technique through which the laboratory chemist can improve processes within a reasonable amount of time compared with traditional conductive synthetic methods. Few examples of chemical and green chemistry-based synthetic routes of hydrazone derivatives are described in **Fig. 8**.

In 2015, Velezheva *et al* have prepared a novel hybrid series of hydrazides and hydrazide-hydrazones possessing the pyrimidine and indole moiety by following **Scheme 1** [64]. In 2017, Popiolek *et al* have reported a novel sequence of hydrazide-hydrazones of isonicotinic acid. The synthetic scheme used by authors is given in **Scheme 2** [57]. In 2017, Zha *et al* have reported a new series of benzo-[d]-thiazole-hydrazone derivatives by using the **Scheme 3** [58]. In 2016, A novel series of N'-substituted-4-methyl benzene sulfonohydrazide derivatives were explored by Ghiya *et al* by using **Scheme 4** [59]. In 2016, A novel class of small molecules of benzothiazole-hydrazones were investigated by Sarkar *et al* using **Scheme 5** [60]. In 2016, Gomathi *et al* have synthesized a novel class of hydrazone schiff's bases. Different hydrazone schiff's bases were prepared by using the following **Scheme 6** [65]. In 2017, Fadare *et al* have investigated a new class of monocarbonyl curcumin analogues and their 2, 4-dinitrophenyl hydrazones using **Scheme 7** [66]. In 2012, Torje *et al* have reported a novel sequence of Phenothiazine-carboxaldehyde-hydrazones and bis-hydrazones through **Scheme 8, 13** [67]. In 2014, Yang *et al* presented a series of novel fluoro-substituted coumarin hydrazones via **Scheme 9** [68]. In 2015, A novel series of mesogenic substituted aroylhydrazones were explored by Singh *et al* through **Scheme 10** [69]. In 2003, Al-Zaydi *et al* described a new series of 2-arylhydrazonopropanals using **Scheme 11** [70]. In 2010, A new series of highly functionalized 4-methyl-1H-quinolin-2-ones were identified by Siddiqui *et al* via **Scheme 12** [71]. In 2011, Aboul-Fadl *et al* identified a new sequence of microwave-assisted one-step synthesis of fenamic acid hydrazides via **Scheme 14** [72].




**Fig. 8.** The graphical representation of some green chemistry-based synthetic routes of hydrazone analogues.

## 4. LITERATURE SURVEY OF HYDRAZONES


### 4.1 Antimicrobial Activity

**Kamal *et al* (2019)**, reported a novel class of 2-arylidene-1-(4-methyl-6-phenylpyrimidine-2-yl) hydrazine analogues and their antimicrobial activity was tested against two Gram-positive bacterial strains (*B. subtilis* MTCC-121 and *S. aureus* MTCC-96) and two Gram-negative bacterial strains (*E. coli* MTCC-1652 and *P. aeruginosa* MTCC-741). Most of the synthesized compounds displayed promising antimicrobial activity against both Gram-positive bacterial strains. The result showed that the compound **1** displayed the potent antimicrobial activity against *B. subtilis* and *S. aureus* with significant MIC values of 50  $\mu\text{g/ml}$  and 100  $\mu\text{g/ml}$ , respectively and the compounds **2**, **3** and **4** showed good activity against *S. cerevisiae* having a zone of inhibition values from 16 mm to 17 mm [73].






**Rezki *et al* (2019)**, reported a novel series of pyridinium hydrazones and phenoxy conjugates. Their antibacterial activity was tested against two strains of Gram-positive bacteria, *Clostridium difficile* and MRSA (Methicillin-Resistant *Staphylococcus aureus*) and two strains of Gram-negative bacteria, *Escherichia coli* and *Neisseria gonorrhea* and their antifungal activity was tested against *Candida albicans*. The result showed that the compound **5** against MRSA, E.C with (MIC=32, 64 µg/ml) and compound **6** (MIC=16 µg/ml) toward MRSA displayed potent antibacterial and antifungal activity [74].




**Khattab *et al* (2019)**, investigated a series of novel hydrazone dyestuff's and screened towards *E. coli*, *S. aureus* and *C. albicans* for their antimicrobial activity. It was determined that analogue **7** displayed the effective antibacterial activity, possessing the bacterial reduction of 10±1.4% against *S. aureus* and 9±1.3% against *E. coli* and antifungal activity towards *C. albicans*, which suggested that these dyes may be used in future for the treatment of the bacterial and fungal infection [63].





**Popiolek *et al* (2018)**, reported a series of novel 2,3-dihalo-substituted propionic acid hydrazides and hydrazide-hydrazones and screened for their *in-vitro* antimicrobial activity

towards Gram-positive strains (*S. aureus* ATCC-25923, *S. aureus* ATCC-43300, *S. aureus* ATCC-6538, *B. subtilis* ATCC-6633, *B. cereus* ATCC-10876, *S. M. luteus* ATCC-10240, *epidermidis* ATCC-12228) and Gram-negative strains (*B. bronchiseptica* ATCC-4617, *E. coli* ATCC-25922, *P. aeruginosa* ATCC-9027, *S. typhimurium* ATCC-14028, *K. pneumoniae* ATCC-13883, *P. mirabilis* ATCC-12453) and antifungal activity against yeast (*C. albicans* ATCC-10231, *C. and Parapsilosis* ATCC-22019). Amongst them, the compound **8** with MIC value of 62.5 µg/ml showed comparable activity as compared to ampicillin with MIC value of 62.5 µg/ml and the compound **9** displayed strong antibacterial activity in response to all bacterial strains, having MIC values in ranges between 31.25-125 µg/ml and MBC values from 62.5-250 µg/ml [57].




**Abdelrahman et al (2017)**, reported two novel series of nicotinic acid hydrazones and quinolone hydrazone analogues. They were tested for their *in-vitro* antimicrobial activity. Results showed that the compounds **10** and **11** bearing 3,4,5-trimethoxy and 2,5-dimethoxy benzylidene moieties were found to be exhibited the excellent antibacterial and antifungal activity, having significant MIC values in the range between 0.49 to 1.95 and 0.49 to 0.98 µg/ml, respectively. Also, several derivatives (**12-23**) displayed significant antimicrobial activities with MIC values in the range between 1.95 to 7.81 µg/ml [62].






**Ajani et al (2017)**, introduced a new series of quinoline based 4-hydrazide hydrazone analogues and examined them for their antibacterial activity towards six bacterial species such as *E. coli*, *S. aureus*, *B. lichenformis*, *M. varians*, *P. aeruginosa* and *P. vulgaris*. Maximum of the analogues exhibited the highest antibacterial activity. It was concluded that compound **24** displayed the most potent antibacterial activity towards all bacterial strains having MIC values in the range between 1.59-0.39 µg/ml [75].




**Cordeiro et al (2016)**, described some isoniazid-derived hydrazone analogues and tested for their *in-vitro* antifungal activity towards *Coccidioides posadasii*. Most of the synthesized derivatives exhibited good antifungal activity but the compound **25** showed the most potent antifungal activity towards *Coccidioides posadasii* strain with MIC value of 100-400 µg/ml. Thus, the results of the present identification revealed that in future, compound **25** may be used against fungal infections [76].




**Kethireddy et al (2015)**, reported a novel series of 5, 6, 7, 8-tetrahydroimidazo [1, 2-a]


pyrimidine-2-carbohydrazide and evaluated for their antibacterial activity towards several bacterial strains. It was revealed that the analogues **26**, **27** and **28** displayed the maximum antibacterial activity against two bacterial strains Gram-positive bacteria (*S. aureus*) and Gram-negative bacteria (*E. coli*) with a zone of inhibition 30-33 mm [77].




**Kaplancıklı et al (2014)**, investigated some novel pyrimidine hydrazone derivatives and evaluated them against several species including, *E. coli*, *B. cereus*, *P. aeruginosa*, *B. subtilis*, *S. typhimurium*, *S. marcescens*, *S. epidermidis* and *C. utilis* for their antimicrobial activity. All the synthesized derivatives displayed potent antimicrobial activity. It was detected that analogue **29** exhibited a strong antimicrobial activity towards all the evaluated species with MIC values in the range between 31.25-250  $\mu\text{g/ml}$  [78].




**Singh et al (2013)**, explored a novel sequence of phenyl hydrazones containing pyrimidine and pyrazole derivatives and tested for their antibacterial activity. Maximum of the prepared derivatives displayed potent antibacterial activity. Among them, compound **30** showed the most promising antibacterial activity towards *B. cereus* and *B. subtilis* with MIC values of 6.25  $\mu\text{g/ml}$  and 1.56  $\mu\text{g/ml}$ , respectively [79].




**Sridhar *et al* (2013)**, scrutinized a novel series of (*E*)-N'-(substituted-benzylidene)-2-(7-fluoro-2-methoxyquinolin-8-yl) acetohydrazide-hydrazone analogues and examined them for their antibacterial activity toward several bacterial strains including, *E. coli* (MTCC-443), *S. aureus* (MTCC-96), *S. pyogenes* (MTCC-442) and *P. aeruginosa* (MTCC-424). Most of the compounds displayed an excellent antibacterial activity. Among them, compounds **31** and **32** exhibited the strong antibacterial activity from 18-21 mm with the inhibition region. SAR study of analogues revealed that the existence of 2, 4-difluoro and 3, 4-difluoro moiety exhibited the maximum antibacterial activity [80].




**Govindaswami *et al* (2011)**, illustrated some novel vanillin related hydrazone derivatives and screened them towards two bacterial strains Gram-positive bacteria (*S. aureus*) and Gram-negative bacteria (*P. aeruginosa*) for their antibacterial activity. Most of these derivatives showed moderate antibacterial activity as compared to the marketable compounds. Among them, the compound **33** was found to be most potent towards *S. aureus* (6-8 mm) and compound **34** showed potent activity towards *P. aeruginosa* (6-12 mm) at 200-250 µg/ml [81].




**Asati *et al* (2011)**, reported a series of novel 1,3-benzothiazole-2-yl-hydrazone analogues and tested for their antibacterial activity towards several bacterial strains (*E. coli*, *B. subtilis*, *P. alkaligenes*, *K. pneumoniae*) and antifungal activity towards fungal strains (*A. niger*, *R. oryzae* and *C. albicans*). Results indicated that compounds **35** and **36** displayed the excellent antibacterial and antifungal activity towards all strains of bacteria and fungus with a zone of inhibition of 18-23 mm [82].




**Ajani *et al* (2010)**, developed a new class of 2-quinoxalinone-3-hydrazone analogues by using the technique of microwave irradiation. They were evaluated for their antimicrobial activity which showed that this molecular context displayed the noticeable effectiveness as antimicrobial agents. The result indicated that the analogue **37** exhibited utmost promising antibacterial activity and compound **38** displayed the most potent antifungal action [83].



**Deep et al (2010)**, invented a sequence of new biphenyl-4-carboxylic acid hydrazide-hydrazoneanalogues and characterized them by using analytical and spectral methods. They were evaluated for their *in-vitro* antimicrobial activity towards two strains of Gram-positive bacteria (*B. subtilis* and *S. aureus*) and two strains of Gram-negative bacteria (*P. aeruginosa* and *E. coli*) and two fungal strains (*A. niger* and *C. albicans*). Most of the analogues displayed effective antimicrobial activity. The result showed that compound **39** exhibited the most potent antibacterial activity, having lower antifungal action and compound **40** exhibited the most potent antifungal activity, having lower antibacterial action [84].



**Khan et al (2009)**, described and investigated some novel hydrazone derivatives of quinoxalinone and characterized them by FT-IR and <sup>1</sup>H-NMR data. Their *in-vitro* antimicrobial activity was evaluated towards two bacterial strains Gram-positive bacteria (*S. aureus* ATCC-29213) and Gram-negative bacteria (*E. coli* ATCC-25922). It was revealed that compounds **41**, **42**, **43** and **44** exhibited the comparatively good activity towards both types of bacterial strains, using ofloxacin as the reference drug. Also, this study showed that the compounds containing extremely electronegative substituents (chloro and fluoro) at the *para*-position of phenyl ring displayed better activity in comparison to the analogues containing these atoms either at ortho or meta-positions or the other analogues comprising fewer electronegative/electropositive substituents at these positions [85].



## 4.2 Antiviral Activity


In the early 20<sup>th</sup> century, infectious diseases were the most common cause of human illness leading to the death [86]. Malignant tumors and viral infections represent the world's most significant threats to public health [87]. Based on current scenario, the spread of Corona Virus Disease (COVID-19) has become a major global public health event, threatens to people's physical and mental health and even life protection, affecting all age groups with a higher incidence in geriatric populations and chronic diseases [88,89]. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is a strain of corona virus that causes coronavirus disease (COVID-19). In these days, spread of SARS-CoV-2 virus through air has become a controversial subject among scientists. Various organizations provide standard methods for monitoring the biological agents [90].

Recently available most of the therapeutically active antiviral medications are intended against human immune deficiency virus (HIV), herpes viruses (best known for causing the cold sores and genital herpes, but causing a wide variety of diseases), hepatitis B and C viruses and influenza A and B viruses [91]. However, prolonged treatment with antiretroviral therapy results in the disease progression. Nowadays, a great advancement requires in the production of newer drugs which may serves as targeted essential aspects of the HIV life cycle in the field of research and development. This includes viral entry (fusion inhibitors) and penetration into the host genome (integrase inhibitors). There is an increasing demand of New Chemical Entities (NCE's) and drug substances due to the increased resistance, recurrence and spread of various diseases, growing number of immunocompromised patients and emergence of many new infections. These have made it possible to identify new medications that are useful in clinical therapy in order to face good health conflict [92]. A large number of hydrazone derivatives have been synthesized and studied for the wide range of antiviral activity.


**Yang *et al* (2018)**, developed a novel series of pyrazole-hydrazone analogues bearing an isoxazole skeleton. They were evaluated for their antiviral activity by using antiviral bioassay(half-leaf method)which indicated that few derivatives showed better *in vivo* antiviral activity towards tobacco mosaic virus (TMV). Amongst, compounds **45**, **46** and **47** displayed the best protection activity, inactivation activity and therapeutic activity against TMV, respectively, which were greater than standard drug Ningnanmycin. This study confirmed that this series of



novel pyrazole-hydrazone analogues containing an isoxazole amide moiety can be considered for the further development as a new class of tobacco protection agents [93].




**Shah *et al* (2018)**, reported a new class of 8-hydroxyquinoline-hydrazone derivatives and characterized them by using  $^1\text{H-NMR}$ ,  $^{13}\text{C-NMR}$ , HRMS and IR techniques. All the synthesized analogues were tested for their anti-HIV-1 activity. Results showed that the compound **48** was found to be most potent ( $\text{IC}_{50}$ : 1.88 and 6.27  $\mu\text{M}$ , TI: 73.82 and 22.07, toward HIV-1<sub>VB59</sub> and HIV-1<sub>UG070</sub>, respectively). This compound **48** (4-methoxy substitution on phenyl ring) exhibited the highest anti-HIV-1 activity (6.5 and 1.6 folds higher) with selectivity (TI approximately 12 and 3 times) against both HIV-1 strains as compared to the parent compound. Thus, the compound **48** can be considered as a lead for improving its anti-HIV-1 potential [30].




**Singh *et al* (2013)**, prepared a novel sequence of phenyl hydrazone comprising pyrazole and pyrimidine derivatives and evaluated against TZM-bl cells for their anti-HIV activity at the concentration of 50  $\mu\text{g ml}^{-1}$ . The result indicated that most of the analogues displayed

promising anti-HIV activity. It was observed that at a test dose, compounds **49**, **50**, **51** and **52** displayed more than 85% inhibition of HIV while analogue **53** showed 98% inhibition. It was demonstrated that these designed molecules have the possibility of introducing the chemical diversity around the core skeleton to generate newer potent molecules [79].




**Ma et al (2011)**, invented a novel sequence of aryl-2-[(4-cyanophenyl)amino]-4-pyrimidinone hydrazone derivatives and tested toward human immunodeficiency virus (HIV)-1 in MT-4 cells for their antiviral activity. Most of the compounds showed tremendous activity towards wild-type HIV-1, with EC<sub>50</sub> values between 1.7-13.2 nm. Among them, the compounds **54** (EC<sub>50</sub>=2.4±0.2nm, SI=18461), **55** (EC<sub>50</sub>=2.6±1.2 nm, SI=2673) and **56** (EC<sub>50</sub>=1.7±0.6 nm, SI=5762) presented the effective activity as compared to ETV toward strains of wild-type HIV-1. Furthermore, the activity of compounds **54** and **56** against HIV-1 RT<sub>(WT)</sub>, CCR5-tropic HIV, and CXCR4-tropic HIV indicates that the target of hyd-CH<sub>2</sub>-DAPYs is wild-type RT only [94].




**Zhan et al (2010)**, reported a novel sequence of N'-arylidene-2-[1-(naphthalen-1-yl)-1H-tetrazol-5-ylthio] acetohydrazide analogues as nonnucleoside reverse transcriptase inhibitors (NNRTIs) and screened them using IIIB strain and ROD strain for their *in vitro* HIV-1 and HIV-2 activity, respectively. The activity was scrutinized via the retardation of virus persuaded cytopathic effects in human T-lymphocyte (MT-4) cells. Results revealed that the analogues **57** and **58** showed EC<sub>50</sub> values of 29.62 μM (CC<sub>50</sub> of 169.24 ± 23.83 μM) and 31.62 μM (CC<sub>50</sub>> 309.06 μM) and resulting in selectivity index of 6 and >9, respectively. The compounds


nevirapine (NVP), efavirenz (EFV), delavirdine (DLV) and zidovudine (azidothymidine, AZT) were used as the reference drugs [95].



**El-Sabbagh *et al* (2009)**, synthesized a novel series of hydrazone derivatives from cyclic  $\beta$ -diketone and their structures were characterized by utilising elemental analysis and numerous spectroscopic techniques. They were screened toward Hepatitis-A Virus (HAV) for their antiviral activity by using the plaque infectivity reduction assay method. It was detected that the compound **59** demonstrated the tremendous antiviral activity, in comparison to the reference drug amantadine [87].




**Vicini *et al* (2009)**, explored a novel sequence benzo[*d*]isothiazolehydrazoneanalogues and examined for their anti-HIV activity in MT-4 cells culture toward wild type strains of HIV and HIV-1 which carry clinically appropriate genetic variations (EFV<sup>R</sup>, Y181C and K103/ Y181C). It was revealed that in this evaluated series, benzo[*d*]isothiazol-3(2*H*)-one moiety was vital for anti-HIV activity. Results showed that analogues **60** and **61** displayed decent activity towards HIV-1 wild type strains, while the analogues **60**, (**62-69**) exhibited good activity towards EFV<sup>R</sup> mutant. It was observed that only one analogue **64** presented noteworthy activity towards Y181C mutant [96].





### 4.3 Anti-TB Activity





**Dogan *et al* (2020)**, developed a novel series of thiadiazolylhydrazones and were tested for their *in vitro* antitubercular activity against *Mycobacterium tuberculosis* H37Rv using Microplate Alamar Blue Assay (MABA) method. Most of the compounds showed good antitubercular activity in the range of MIC 0.78-6.25 µg/ml. Compounds **70** and **71** showed prominent antitubercular activity at MIC of 0.78 µg/ml. It is concluded that these compounds possessed antitubercular activity due to enoyl acyl carrier protein reductase (InhA) as the possible target enzyme [97].



**Sutar *et al* (2019)**, prepared a new series of isoniazid-transferrin conjugates viz TF-IH and TF-IG (**72**, **73**). They were further investigated for their *in-vitro* anti-tubercular activity. These conjugates were found to have higher pH-dependent stability and better anti-TB activity than free isoniazid. Transferrin-IH conjugates exhibited a significant drop in the colony-forming unit at 0.39 µg/ml conc. than Transferrin-IG & free INH during *ex-vivo* studies. Hence, it's evident that the conjugation of anti-tuberculosis agents with transferrin improves the potency and efficiency of drugs [98].




**Sampiron *et al* (2019)**, presented a novel sequence of hydrazone, benzohydrazones and isoniazid-acylhydrazones derivatives and determined for their *in vitro* antitubercular activity against *M. tuberculosis* H<sub>37</sub>Rv and other various strains. All compounds revealed decent activity at MIC of 0.12-250 µg/ml. They exhibited significant MIC at both pH 6.8 and 6.0. Compounds **74-77** with isoniazid-acylhydrazones showed prominent results. Based on the above results, isoniazid-acylhydrazones can be used for new drug development as antitubercular agents [99].




**Hakkimane *et al* (2018)**, reported a novel sequence of rifampicin-loaded poly lactic-co-glycolic acid nanoparticles and isoniazid derived isoniazid benz-hydrazones and tested for their antitubercular activity against the H37Rv strain of *Mycobacterium tuberculosis*. It was observed that


analogue **78** showed potent antimycobacterial activity, better stability and sustain release properties for one month. All the synthesized compounds were found to be better anti-TB agents and have more efficacy in drug delivery than free forms [100].




**Bonnett et al (2018)**, explored a series of hydrazones and examined for their antimycobacterial activity toward non-replicating *M. tuberculosis* species. All the parameters like Minimum Inhibitory Concentration (MIC), Low Oxygen Recovery Assay (LORA) and Minimum Bactericidal Concentration (MBC) were recorded for tested compounds. Most of them showed excellent anti-tubercular activity. Amongst, two compounds **79** and **80** exhibited the remarkable activity under hypoxic conditions than aerobic culture. In a nutshell, hydrazones have prominent activity as antitubercular agents [101].



**Luet al (2017)**, reported two novel series of benzylsulfanyl benzo-heterocyclic amides and hydrazones derivatives. These compounds were assessed for their anti-TB activities and some of the isonicotinylhydrazones derivatives possessed decent anti-tubercular activity towards the H37Rv strain of *M. tuberculosis*. MIC values of these compounds were found to be better than isoniazid, moxifloxacin, streptomycin and PA-824. Further, *in-vivo* studies explored that compound **81** can be used as a new anti-TB agent as it significantly lowers the activity of mycobacterium in H37Ra infected mouse model [102].




**Abdelrahman et al (2017)**, have identified a new sequence of pyridine and quinoline based hydrazone analogues. The prepared analogues were evaluated for their anti-TB activity. Results concluded that analogues **82**, **83** and **84** displayed remarkable antimycobacterial (MIC = 0.39 µg/ml). Whereas, the compounds **85**, **86**, **87** and **88** displayed excellent antimycobacterial activity (MIC = 0.78 µg/ml) [62].




Comp	R	R <sub>1</sub>
<b>82</b>	4-Cl	2-OH-3-OCH <sub>3</sub>
<b>83</b>	4-Cl	2,5-(OCH <sub>3</sub> ) <sub>2</sub>
<b>84</b>	3-Cl	2,5-(OCH <sub>3</sub> ) <sub>2</sub>
<b>85</b>	3-Cl	3,4-(OCH <sub>3</sub> ) <sub>2</sub>
<b>86</b>	3-Cl	3,4,5-(OCH <sub>3</sub> ) <sub>3</sub>
<b>87</b>	4-Cl	3,4,5-(OCH <sub>3</sub> ) <sub>3</sub>
<b>88</b>	4-Cl	4-OCH <sub>3</sub>


**Joshi et al (2016)**, prepared a novel sequence of pyrrolyl hydrazones and their Cu(II) complexes. The compounds were analyzed *in-vitro* for their anti-tubercular activity against the bacterial strains of *M. tuberculosis*. As compared to ligands metal-complexes were found to be more potent. Compounds (**89-91**) showed appreciable anti-TB activity with MIC value of 0.8 µg/ml against standard drug rifampicin. These compounds were able to bind more than 60% with enzymes at a concentration lower than 5 µM [103].




**Cihan-Ustunda<sup>†</sup> et al (2015)**, developed a novel series of indoyl hydrazones and indole-based 4-thiazolidinones. The synthesized analogues were examined *in-vitro* for their anti-tubercular activity towards *Mycobacterium tuberculosis* H37Rv. It was found that compounds **92-96** showed excellent anti-TB activity with a 10-fold lower concentration of mammalian cell toxicity with MIC value of 6.25 to 25.0 µg/ml [104].









**Abdel-Aziz et al (2015)**, presented three novel series of halophenyl bis-hydrazones and tested for their antimycobacterial activities. Results indicated that analogues **97-101** possessing broad antimicrobial activity against *Mycobacterium tuberculosis*. The above mentioned five analogues can serve fairly well to recognize chemotherapeutics of broad-spectrum anti-TB [105].






**Dandawate *et al* (2014)**, investigated a new sequence of four analogues (**102-105**) of ferrocenylhydrazones under varied iron conditions and their  $\beta$ -cyclodextrin (CD) inclusion complexes were synthesized and examined for their anti-tubercular activity. It was discovered that low MIC values (0.5-128  $\mu\text{g/ml}$ ) against 7H9 medium were shown by cyclodextrin inclusion complex of INH. FBHZ and its cyclodextrin inclusion complex was found to be the most significant amongst them and exhibited MIC values of 1.0 and 0.5  $\mu\text{g/ml}$ . Cyclodextrin conjugates hold good potential for the production of antitubercular drugs under varied iron condition [106].




**Jamadaret *al* (2012)**, arranged a new sequence of pyruvate based hydrazones and tested toward *M. tuberculosis* using middle brook 7H9 medium. About eight of the pyruvate hydrazones and its sodium salts displayed inactivity in their anionic form. On the other side, prominent antimycobacterial activity was displayed by their neutral Cu (II) complexes (**106-113**) under elevated iron (8  $\mu\text{g Fe per ml}$ ) condition. The complexes  $[\text{Cu}(\text{L}^1)(\text{H}_2\text{O})_2] - [\text{Cu}(\text{L}^8)(\text{H}_2\text{O})_2]$  showed MIC values of <0.5 - 2  $\mu\text{g/ml}$  [107].




Comp	X	R	Comp	X	R
<b>106</b>	CH	4-CH <sub>3</sub>	<b>110</b>	CH	4-CH <sub>3</sub>
<b>107</b>	CH	4-NO <sub>2</sub>	<b>111</b>	CH	3-F
<b>108</b>	CH	4-(CH <sub>3</sub> ) <sub>3</sub>	<b>112</b>	CH	3-OH
<b>109</b>	CH	4-NO <sub>2</sub> ,3-CH <sub>3</sub>	<b>113</b>	N	H


**Mahajan *et al* (2011)**, introduced a novel series of ferrocene based hydrazones and evaluated for their anti-tuberculosis activity towards *Mycobacterium tuberculosis*. Among the tested analogues,analogue **114** exhibited potent anti-TB activity against *M. tuberculosis*, having a MIC value of 2.5-5 µg/ml. Results revealed that such hybrid analogues provide an efficient approach for future pharmacological developments to fight against tuberculosis [21].




**Pinheiro *et al* (2011)**, explored a series of *L*-serniylhydrazone derivatives and screened for their *in-vitro* antitubercular activity towards *Mycobacterium tuberculosis* H37Rv. Generally,the tested analogues were tested for antitubercular activity and showed good activity. The compounds **115-120** showed remarkable activity in the range of MIC 25 and 100 µg/ml, and were comparable with standard antitubercular drug D-cicloserine (5-20 µg/ml) [108].




Comp	R <sub>1</sub>	R <sub>2</sub>	R <sub>3</sub>
<b>115</b>	H	NO <sub>2</sub>	H
<b>116</b>	H	CN	H
<b>117</b>	OH	OH	H
<b>118</b>	OH	H	OH
<b>119</b>	OH	OCH <sub>3</sub>	H



**Kumar *et al* (2010)**, identified a novel series of benzoic acid hydrazones and their nicotinyl derivatives. The synthesized analogues were assayed *in-vitro* for their anti-tubercular activities against *Mycobacterium tuberculosis* strain. It was revealed that compound **121** showed the most potent anti-tubercular activity with MIC value of  $3.5 \times 10^{-3} \mu\text{M}$  [109].



**Bairwa *et al* (2010)**, reported the synthesis of a novel sequence of phenylacrylamide analogues substituting cinnamic acid and guanylhydrazones by employing microwave-assisted synthesis. Resazurin-microtitre plate assay (REMA) was used to evaluate the anti-tubercular activity of compounds against *Mycobacterium tuberculosis*. Results demonstrated that compound **122** is of particular interest having good safety profile with MIC of  $6.48 \mu\text{M}$ . In future, the compound can be used for assisting antitubercular studies [110].



## 5. RECENT PATENTS FILED/GRANTED ON HYDRAZONE DERIVATIVES

S. No.	Date of Publications	Patent Number	Inventions Disclosed	Reference s
1.	07/11/2017	US 9,809,550 B2	Quinolyl hydrazones for the treatment of tuberculosis and related diseases.	[111]
2.	14/03/2017	US 9,593,103 B2	Pyridyl hydrazones for the treatment of tuberculosis and related diseases.	[112]
3.	15/09/2015	US 9,133,230 B2	Hydrazone derivatives having potent antitumor activity toward multi-drug resistant tumor cells.	[113]
4.	07/01/2014	US 8,623,849 B2	Medicinal applications of benzoic acid hydrazones synthesized on the basis of steroidal tigogenin.	[114]
5.	30/07/2013	US 8,497,296 B2	N, N'-hydrazino-bis-isatin derivatives with selective activity against multidrug-resistant cancer cells.	[115]
6.	24/01/2012	US 8,101,649 B2	N-acyl hydrazone derivatives useful as modulators of nicotinic acetylcholine receptors.	[116]
7.	06/04/2010	US 7,692,030 B2	Conjugates of artemisinin-related endoperoxides and hydrazone derivatives for the treatment of cancer.	[117]
8.	20/08/2009	US2009/020837 4 A1	Novel silane compounds carrying a hydrazone or diazo functional group in order to functionalize solid supports and immobilize biological molecules on these supports.	[118]
9.	01/04/2008	US 7,351,831 B2	Process for asymmetric intramolecular (3+2) cyclo-addition of hydrazones.	[119]
10.	11/10/2007	US 2007/0238700	N-phenyl-1,1,1-trifluoromethane-sulfonamide hydrazone compounds and	[120]

		A1	their usage in controlling parasites.	
11.	07/12/2006	US 2006/0276433 A1	Hydrazone derivatives.	[121]
12.	09/12/2003	US 6,660,737 B2	Medicinal uses of hydrazones.	[122]
13.	18/03/2003	US 6,534,502 B2	Substituted oximes and hydrazones as neurokinin antagonists.	[123]

## 6. CONCLUSION

The invention of novel chemotherapeutic agents is an essential and challenging task for the medicinal chemists and several research programs are directed towards the design and synthesis of newer drugs for their chemotherapeutic usage. To discover active analogues towards multidrug-resistant microbial infections, hydrazone containing analogues are found as an important class for the development of newer anti-infective drugs. Hydrazone ligands attracted the special attention of researchers due to their well-known chelating capability, structural flexibility and diverse range of pharmaceutical applications. The present review highlights the potential of hydrazone derivatives as a guideline for the advancement of anti-infective agents, SAR based study of hydrazone analogues with a diverse mechanism of action of nanoparticles based hydrazone compounds as potent anti-infective agents have been addressed in effective way. Various conventional and green synthesis of hydrazone derivatives and recent patents filed or granted on hydrazone compounds as anti-infective agents have also been tabulated. It could be observed that with careful designing of known hydrazone derivatives and future work might yield more promising lead compounds with better anti-infective activity.

## 7. FUTURE ASPECTS:

Hydrazone is indeed one of the primary drug research scaffolds due to its remarkable therapeutic properties. In heterocyclic chemistry, the expansion of effective and reliable methods for their synthesis and fusion with many other bioactive molecules has ensured that hydrazones have been a crucial moiety. Research thresholds will therefore remain crucial to the discovery of bioactive

molecules in order to satisfy all the relevant criteria for the expansion of potent antiviral and antimicrobial drugs. Current and re-emerging infectious diseases will continue to present significant challenges to global health into the 21<sup>st</sup> century. According to world health organization (WHO) study, these are already considered as leading cause of death and affected patient health care system across the world. Because of the rising number of infections and resistance towards conventional treatments, there is an persistent need to produce some useful medications that can counter resistance and side-effects related to anti-infective medications. In addition, rational design and production of the novel anti-infective agents that contain this scaffold will help to resolve increasing microbial resistance problems and satisfy the need for successful antimicrobial therapy to combat multiple deadly microbial infectious diseases. Based on the literature survey, we ascertained that the future of medicinal chemistry should focuses on the development of newer molecules with innovative approach in order to obtain therapeutically active molecules in forthcoming future. The development of such new molecules includes new synthetic instruments, synthetic strategies and techniques.

#### ACKNOWLEDGEMENT

The authors PCS and VKT would like to express their sincere thanks to Newton Bhabha Fund for providing the foundation to establish this cross-country interdisciplinary research collaboration through Researcher Links scheme.

#### CONFLICT OF INTEREST

There are no conflicts of interest in this article.

#### ABBREVIATIONS

HAV	Hepatitis-A Virus	IR	Infrared
ATCC	American Type Culture Collection	NMR	Nuclear magnetic resonance
DNA	Deoxyribonucleic Acid	SAR	Structure-activity relationship
MRSA	Methicillin-resistant Staphylococcus aureus	REMA	Resazurin Microtiter Assay Plate
MAOS	Microwave-assisted organic synthesis	MTCC	Microbial Type Culture Collection

TMV	Tobacco mosaic virus	MIC	Minimum inhibitory concentration
μM	Micromolar	MBC	Minimum bactericidal concentration
Mm	Millimeter	ETV	Entecavir
Nm	Nanometer	LORA	Low Oxygen Recovery Assay
μg/ml	Microgram per millilitre	FBHZ	Ferrocenylhydrazone
MABA	Microplate Alamar Blue Assay	ROS	Reactive oxygen species
HRMS	High Resolution Mass Spectrometry	TZM- <i>bl</i>	Hela cell line
HIV	Human Immunodeficiency Virus	EC <sub>50</sub>	Half maximal effective concentration
SI	Standard international	INH	Isoniazid
ROD	Varicella-Zoster Virus strain		

## REFERENCES

- [1] R. Bhaskar, N. Salunkhe, A. Yaul, A. Aswar, Bivalent transition metal complexes of ONO donor hydrazone ligand: Synthesis, structural characterization and antimicrobial activity, *Spectrochim. Acta A Mol. Biomol. Spectrosc.* 151 (2015) 621-627.
- [2] R. Fekri, M. Salehi, A. Asadi, M. Kubicki, Synthesis, characterization, anticancer and antibacterial evaluation of Schiff base ligands derived from hydrazone and their transition metal complexes, *Inorg. Chim. Acta* 484 (2019) 245-254. <https://doi.org/10.1016/j.ica.2018.09.022.0>.
- [3] A.S. Burlov, V.G. Vlasenko, Y.V. Koshchienko, N.I. Makarova, A.A. Zubenko, Y.D. Drobin, L.N. Fetisov, A.A. Kolodina, Y.V. Zubavichus, A.L. Trigub, Synthesis, characterization, luminescent properties and biological activities of zinc complexes with bidentate azomethine Schiff base ligands, *Polyhedron* 154 (2018) 65-76.
- [4] D.L. Peng, Synthesis, crystal structures and catalytic properties of two oxido-vanadium (V) complexes with tridentate Schiff bases, *Russ. J. Coord. Chem.* 43(6) (2017) 404-410.
- [5] D.L. Peng, Crystal structure and catalytic property of an oxido-molybdenum (VI) complex derived from N'-(2-hydroxy-3, 5-di-tert-butylbenzylidene)-4-methylbenzo hydrazide, *J. Struct. Chem.* 59(3) (2018) 589-594.
- [6] G. Gurkok, N. Altanlar, S. Suzen, Investigation of antimicrobial activities of indole-3-



- aldehyde hydrazone/hydrazone derivatives, *Chemotherapy*. 55(1) (2009) 15-19.
- [7] G. Uppal, S. Bala, S. Kamboj, M. Saini, Therapeutic review exploring antimicrobial potential of hydrazones as promising lead, *Der Pharma Chem*. 3(1) (2011) 250-268.
- [8] S. Rollas, S.G. Kucukguzel, Biological activities of hydrazone derivatives, *Molecules*. 12(8) (2007) 1910-1939.
- [9] L. Wang, D.G. Guo, Y.Y. Wang, C.Z. Zheng, 4-Hydroxy-3-methoxy-benzaldehyde series aroylhydrazones: Synthesis, thermostability and antimicrobial activities, *RSC Adv*. 4(102)(2014) 58895-58901.
- [10] D.L. Peng, N. Sun, Synthesis, characterization and crystal structures of Schiff base zinc (II) complexes with antibacterial activity, *Acta Chim. Slov*. 65(4) (2018) 895-901.
- [11] H.Y. Qian, Synthesis, crystal structures and biological activity of manganese (III) complexes derived from bis-Schiff bases, *Russ. J. Coord. Chem*. 44(1) (2018) 32-38.
- [12] N. Ribeiro N, S. Roy, N. Butenko, I. Cavaco, T. Pinheiro, I. Alho, F. Marques, F. Avecilla, J.C. Pessoa, I. Correia, New Cu (II) complexes with pyrazolyl derived Schiff base ligands: Synthesis and biological evaluation, *J. Inorg. Biochem*. 174 (2017) 63-75.
- [13] L. Mitu, M. Ilis, N. Raman, M. Imran, S. Ravichandran, Transition metal complexes of isonicotinoyl-hydrazone-4-diphenylaminobenzaldehyde: Synthesis, characterization and antimicrobial studies, *J. Chem*. 9(1) (2012) 365-372.
- [14] M. La, P.P. Wang, L.W. Xue, Synthesis, characterization, crystal structures and antimicrobial activity of oxido vanadium (V) complexes with hydrazone and pyrone ligands, *Inorg. Nano-Met. Chem*. 50(3) (2020) 151-155. <https://doi.org/10.1080/24701556.2019.1694038>.
- [15] M. Alam, G. Verma, M. Shaquiquzzaman, A. Marella, M. Akhtar, M. Ali, A review exploring biological activities of hydrazones, *J. Pharm. Bioallied Sci*. 6(2) (2014) 69.
- [16] Y. Zhao, J. Hui, D. Wang, L. Zhu, J.H. Fang, X.D. Zhao, Synthesis, cytotoxicity and pro-apoptosis of novel benzoisoindolinhydrazones as anticancer agents, *Chem. Pharm. Bull*. 58(10) (2010) 1324-1327.
- [17] T. Nasr, S. Bondock, M. Youns, Anticancer activity of new coumarin substituted hydrazone-hydrazone derivatives, *Eur. J. Med. Chem*. 76 (2014) 539-548.
- [18] R.R. Khattab, A.A. Hassan, O.M. Kutkat, K.M. Abuzeid, N.A. Hassan, Synthesis and antiviral activity of novel thieno-[2,3-d] pyrimidine hydrazones and their C-

- nucleosides, Russ. J. Gen. Chem. 89(8) (2019) 1707-1717.
- [19] Z. Yang, P. Li, X. Gan, Novel pyrazole-hydrazone derivatives containing an isoxazole moiety: Design, synthesis and antiviral activity, *Molecules*. 23(7) (2018) 1798.
- [20] J. Patole, U. Sandbhor, S. Padhye, D.N. Deobagkar, C.E. Anson, A. Powell, Structural chemistry and *in-vitro* antitubercular activity of acetylpyridine benzoyl hydrazone and its copper complex against *Mycobacterium smegmatis*, *Bioorg. Med. Chem. Lett.* 13(1) (2003) 51-55.
- [21] A. Mahajan, L. Kremer, S. Louw, Y. Gueradel, K. Chibale, C. Biot, Synthesis and *in-vitro* antitubercular activity of ferrocene-based hydrazones, *Bioorg. Med. Chem. Lett.* 21(10) (2011) 2866-2868.
- [22] O. Oniga, J.T. Ndongo, C. Moldovan, B.R. Tiperciuc, S.M. Oniga, A.D. Pirnau, L.A. Vlase, P.H. Verite, Synthesis and antimicrobial activity of some new 2-hydrazone-thiazoline-4-ones, *Farmacia*. 60 (2012) 785.
- [23] N. Thilagavathi, A. Manimaran, N. Padma Priya, N. Sathya, C. Jayabalakrishnan, Synthesis, characterization, electrochemical, catalytic and antimicrobial activity studies of hydrazone Schiff base ruthenium (II) complexes, *Appl. Organomet. Chem.* 24(4) (2010) 301-307.
- [24] A. Kajal, S. Bala, N. Sharma, S. Kamboj, V. Saini, Therapeutic potential of hydrazones as anti-inflammatory agents, *Int. J. Med. Chem.* (2014) 1-11.
- [25] A.K. Reddy, N.E. Kathale, Synthesis, characterization and anti-inflammatory activity of hydrazones bearing 5-nitro-furan moiety and 5-iodo-vanillin hybrid, *Orient. J. Chem.* 33(2) (2017) 971-978.
- [26] J.V. Ragavendran, D. Sriram, S.K. Patel, I.V. Reddy, N. Bharathwajan, J. Stables, P. Yogeeswari, Design and synthesis of anticonvulsants from a combined phthalimide-GABA-anilide and hydrazone pharmacophore, *Eur. J. Med. Chem.* 42(2) (2007) 146-151.
- [27] J. Jain, Y. Kumar, R. Sinha, R. Kumar, J. Stables, Menthone aryl acid hydrazones: A new class of anticonvulsants, *Med. Chem.* 7(1) (2011) 56-61.
- [28] A. Inam, S.M. Siddiqui, T.S. Macedo, D.R. Moreira, A.C. Leite, M.B. Soares, A. Azam, Design, synthesis and biological evaluation of 3-[4-(7-chloro-quinolin-4-yl)-piperazin-1-yl]-propionic acid hydrazones as antiprotozoal agents, *Eur. J. Med. Chem.* 75 (2014) 67-76.

- [29] J.C. Coa, W. Castrillon, W. Cardona, M. Carda, V. Ospina, J.A. Munoz, I.D. Velez, S.M. Robledo, Synthesis, leishmanicidal, trypanocidal and cytotoxic activity of quinoline-hydrazonehybrids, *Eur. J. Med. Chem.* 101 (2015) 746-753.
- [30] P. Shah, L.F. Abadi, S. Gaikwad, D. Chaudhari, V. Kushwah, S. Jain, K.K. Bhutani, S. Kulkarni, I.P. Singh, Synthesis and biological evaluation of 8- hydroxyquinoline- hydrazones for anti- HIV- 1 and anticancer potential, *Chemistry Select.* 3(38) (2018) 10727-10731.
- [31] X.D. Ma, S.Q. Yang, S.X. Gu, Q.Q. He, F.E. Chen, E. De Clercq, J. Balzarini, C. Pannecouque, Synthesis and anti- HIV activity of Aryl- 2- [(4- cyanophenyl)- amino]- 4- pyrimidinone hydrazones as potent non- nucleoside reverse transcriptase inhibitors, *Chem. Med. Chem.* 6(12) (2011) 2225-2232.
- [32] N. Belkheiri, B. Bouguerne, F. Bedos-Belval, H. Duran, C. Bernis, R. Salvayre, A. Negre-Salvayre, M. Baltas, Synthesis and antioxidant activity evaluation of a syringic hydrazones family, *Eur. J. Med. Chem.* 45(7) (2010) 3019-3026.
- [33] S.M. Sondhi, M. Dinodia, A. Kumar, Synthesis, anti-inflammatory and analgesic activity evaluation of some amidine and hydrazone derivatives, *Bioorg. Med. Chem.* 14(13) (2006) 4657-4663.
- [34] G. Rajitha, K.V. Prasad, A. Umamaheswari, D. Pradhan, K. Bharathi, Synthesis, biological evaluation and molecular docking studies of N-( $\alpha$ -acetamido cinnamoyl) aryl hydrazone derivatives as anti-inflammatory and analgesic agents, *Med. Chem. Res.* 23(12) (2014) 5204-5214.
- [35] B. Ates, S. Koytepe, A. Ulu, C. Gurses, V. K. Thakur, Chemistry, Structures, and Advanced Applications of Nanocomposites from Biorenewable Resources, *Chemical Reviews* (2020) 10.1021/acs.chemrev.9b00553.
- [36] P.K. Rai, Z. Usmani, V.K. Thakur, V.K. Gupta, Y.K. Mishra, Tackling COVID-19 Pandemic through Nanocoatings: Confront and Exactitude, *Current Research in Green and Sustainable Chemistry*, 2020, 100011.
- [37] S. Thakur, R.V. Saini, P. Singh, P. Raizada, V.K. Thakur, A.K. Saini, Nanoparticles as an emerging tool to alter the gene expression: Preparation and conjugation methods, *Materials Today Chemistry* 17, 2020, 100295.
- [38] J.J. Giner-Casares, M. Henriksen-Lacey, M. Coronado-Puchau, L.M. Liz-Marzán,

Inorganic nanoparticles for biomedicine: where materials scientists meet medical research, *Mater.Today*. 19(1) (2016) 19-28.

- [39] A. Abdel-Azeem, A.A. Nada, A. O'donovan, V.K. Thakur, A. Elkelish, Mycogenic Silver Nanoparticles From Endophytic *Trichoderma atroviride* with Antimicrobial Activity, *Journal of Renewable Materials* 8 (2), 2020, 171.
- [40] P.J. Espitia, N.D. Soares, J.S. dos Reis Coimbra, N.J. de Andrade, R.S. Cruz, E.A. Medeiros, Zinc oxide nanoparticles: Synthesis, antimicrobial activity and food packaging applications, *Food Bioprocess Tech.* 5(5) (2012) 1447-1464.
- [41] V.K. Pandey, K.R. Srivastava, G. Ajmal, V.K. Thakur, V.K. Gupta, S.N. Upadhyay, P.K. Mishra, Differential susceptibility of catheter biomaterials to biofilm-associated infections and their remedy by drug-encapsulated Eudragit RL100 nanoparticles, *International journal of molecular sciences* 20 (20), (2019) 5110.
- [42] M. Arakha, S. Pal, D. Samantarrai, T.K. Panigrahi, B.C. Mallick, K. Pramanik, B. Mallick, S. Jha, Antimicrobial activity of iron oxide nanoparticle upon modulation of nanoparticle-bacteria interface, *Sci. Rep.* 5 (2015) 14813.
- [43] M. Rai, S.D. Deshmukh, A.P. Ingle, I.R. Gupta, M. Galdiero, S. Galdiero, Metal nanoparticles: The protective nanoshield against virus infection, *Crit. Rev. Microbiol.* 42(1) (2016) 46-56.
- [44] R. Singh, Nawale L, Arkile M, Wadhwani S, Shedbalkar U, Chopade S, Sarkar D, Chopade BA. Phytogenic silver, gold, and bimetallic nanoparticles as novel antitubercular agents. *International journal of nanomedicine*. 11 (2016) 1889-1897.
- [45] S. Rahman, L. Rahman, A.T. Khalil, N. Ali, D. Zia, M. Ali, Z.K. Shinwari, Endophyte-mediated synthesis of silver nanoparticles and their biological applications, *Appl. Microbial. Biotechnol.* 103(6) (2019) 2551-69.
- [46] N. Srividya, M.D. Ghora, P.R. Padmanabh, Antimicrobial nanotechnology: research implications and prospects in food safety, In *Food Preservation 2017* (pp. 125-165). Academic Press.
- [47] C.W. Mandl, Steps of the tick-borne encephalitis virus replication cycle that affect neuropathogenesis, *Virus Res.* 111(2) (2005) 161-174.

- [48] R. Pangestika, R. Ernawati, Antiviral activity effect of silver nanoparticles (AgNps) solution against the growth of infectious bursal disease virus on embryonated chicken eggs with Elisa test, *KnE Life Sci.* (2017) 536-548.
- [49] H.B. Rode, D.M. Lade, R. Grée, P.S. Mainkar, S. Chandrasekhar, Strategies towards the synthesis of anti-tuberculosis drugs, *Org. Biomol. Chem.* 17(22) (2019) 5428-5459.
- [50] K. Patil, S. Bagade, S. Bonde, S. Sharma, G. Saraogi, Recent therapeutic approaches for the management of tuberculosis: challenges and opportunities, *Biomed. Pharmacother.* 99 (2018) 735-745.
- [51] J. Wu, S. Kang, B. Song, D. Hu, M. He, L. Jin, S. Yang, Synthesis and antibacterial activity against *Ralstonia solanacearum* for novel hydrazone derivatives containing a pyridine moiety, *Chem. Cent. J.* 6 (2012) 28.
- [52] A.S. Hassan, A.A. Askar, E.S. Nossier, A.M. Naglah, G.O. Moustafa, M.A. Al-Omar, Antibacterial evaluation, *in silico* characters and molecular docking of Schiff bases derived from 5-aminopyrazoles, *Molecules*. 17 (2019) 3130.
- [53] I.A. Khodja, H. Boulebd, Synthesis, biological evaluation, theoretical investigations, docking study and ADME parameters of some 1, 4-bisphenylhydrazone derivatives as potent antioxidant agents and acetylcholinesterase inhibitors, *Mol. Divers.* (2020) 1-2.
- [54] W.M. Eldehna, M. Fares, M.M. Abdel-Aziz, H.A. Abdel-Aziz, Design, synthesis and antitubercular activity of certain nicotinic acid hydrazides, *Molecules*. 5 (2015) 8800-15.
- [55] [www.omicsonline.org>structure-activity-relationship-scholarly-open](http://www.omicsonline.org>structure-activity-relationship-scholarly-open). (Accessed on 04-june-2020)
- [56] E.M. Afsah, E. Abdel-Galil, I. Youssef, Y.H. Elsaedy, Synthesis and biological activity of novel hydrazide-hydrazones incorporating piperidine and piperazine moieties and related mannich bases, *Am. J. Mater. Sci.* 6(2) (2019) 58-65.
- [57] L. Popiolek, A. Biernasiuk, New hydrazides and hydrazide-hydrazones of 2, 3-dihalogen substituted propionic acids: Synthesis and *in-vitro* antimicrobial activity evaluation, *Chem. Biodivers.* 14(8) (2017) e1700075.
- [58] G. Zha, J. Leng, N. Darshini, T. Shubhavathi, H.K. Vivek, A.M. Asiri, H.M. Marwani, K.P. Rakesh, N. Mallesha, H. Qin, Synthesis, SAR and molecular docking studies of benzo-[d]-thiazole-hydrazones as potential antibacterial and antifungal agents, *Bioorg. Med. Chem. Lett.* 27 (2017) 3148-3155.

- [59] S. Ghiya, Y.C. Joshi, Synthesis and antimicrobial evaluation of hydrazones derived from 4-methylbenzene sulfonohydrazide in aqueous medium, *Med. Chem. Res.* 25(5) (2016) 970-976.
- [60] S. Sarkar, A.A. Siddiqui, S.J. Saha, R. De, S. Mazumder, C. Banerjee, M.S. Iqbal, S. Nag, S. Adhikari, U. Bandyopadhyay, Antimalarial activity of small molecule benzothiazole hydrazones, *Antimicrob. Agents Chemother.* 60(7) (2016) 4217-4228. <https://doi.org/10.1128/AAC.01575-15>.
- [61] R. Anbazhagan, K.R. Sankaran, Design, synthesis, computational calculation and biological evaluation of some novel 2-thiazolyl hydrazones, *Spectrochim. Acta Part A. Mol. Biomol. Spectrosc.* 135 (2015) 984-993.
- [62] M.A. Abdelrahman, I. Salama, M.S. Gomaa, M.M. Elasser, M.M. Abdel-Aziz, D.H. Soliman, Design, synthesis and 2D QSAR study of novel pyridine and quinolone hydrazone derivatives as potential antimicrobial and antitubercular agents, *Eur. J. Med. Chem.* 138 (2017) 698-714. <https://doi.org/10.1016/j.ejmech.2017.07.004>.
- [63] T.A. Khattab, A.A. Allam, S.I. Othman, M. Bin-jumah, H.M. Al-harbi, M.M.G. Fouda, Synthesis, solvatochromic performance, pH sensing, dyeing ability and antimicrobial activity of novel hydrazone dyestuffs, *J. Chem.* (2019). <https://doi.org/10.1155/2019/7814179>.
- [64] V. Velezhevaa, P. Brennanb, P. Ivanova, A. Kornienkoa, S. Lyubimova, K. Kazarian, B. Nikonenko, K. Majoro, Synthesis and antituberculosis activity of indole-pyridine derived hydrazides, hydrazide-hydrazones and thiosemicarbazones, *Bioorg. Med. Chem. Lett.* 26(3) (2016) 978-985. <https://doi.org/10.1016/j.bmcl.2015.12.049>.
- [65] G. Gomathi, R. Gopalakrishnan, A hydrazone Schiff base single crystal (E)-methyl physicochemical, *in-vitro* investigation of antimicrobial activities and molecular docking with DNA gyrase protein, *Mater. Sci. Eng. C.* 64 (2016) 133-138.
- [66] O.A. Fadare, E.O. Iwalewa, C.A. Obafemi, F.P. Olatunji, In-silico antimalarial study of monocarbonyl curcumin analogs and their 2, 4-dinitro phenylhydrazones using the inhibition of plasmeprin-II as test model, *JPS.* 5(2) (2017) 18-24.
- [67] I.A. Torje, A.M. Valean, C. Cristea, Phenothiazinecarboxaldehyde-hydrazone derivatives synthesis, characterization and electronic properties, *Rev. Roum. Chim.* 57(2012)337-44.
- [68] G.Y. Yang, J.T. Yang, C.X. Wang, S.F. Fan, P.H. Xie, C.L. Xu, Microwave-assisted TsOH/SiO<sub>2</sub>-catalyzed one-pot synthesis of novel fluoro-substituted coumarin hydrazones

- under solvent-free conditions, *J. Fluor. Chem.* 168 (2014) 1-8.
- [69] S.K. Singh, V. Kumar, H.K. Singh, P.Kanth, B. Singh, Microwave-assisted synthesis, characterisation and mesomorphic investigations of novel disubstituted aroylhydrazones. *Liq. Cryst.*42(8)(2015)1179-90.
- [70] K.M. Al-Zaydi, R.M.Borik, M.H. Elnagdi, 2-Arylhydrazonopropanals as building blocks in heterocyclic chemistry: Microwave assisted condensation of 2-arylhydrazonopropanals with amines and active methylene reagents, *Molecules.* (12) (2003) 910-23.
- [71] I.R. Siddiqui, S. Shamim, A. Singh, V. Srivastava, S. Yadav, Moisture compatible and recyclable indium (III) chloride catalyzed and microwave assisted efficient route to substituted 1H-quinolin-2-ones, *ARKIVOC.* 11 (2010) 232-241.
- [72] T.Aboul-Fadl, H.A. Abdel-Aziz, A.Kadi, A. Bari, P. Ahmad, T. Al-Samani, S.W. Ng, Microwave-assisted one-step synthesis of fenamic acid hydrazides from the corresponding acids, *Molecules.* (5) (2011) 163544-51.
- [73] R. Kamal, R. Kumar, V. Kumar, V. Kumar, K.K. Bansal, P.C. Sharma, Synthesis, anthelmintic and antimicrobial evaluation of new 2-arylidene-1-(4-methyl-6-phenylpyrimidin-2-yl) hydrazines, *J. Med. Chem. Drug Discov.*4 (2019)713-717.
- [74] N.Rezki, S.A. Al-Sodies, H.E.A. Ahmed, S.Ihmaid, M.Messali, S. Ahmed, M.R.Aouad, A novel dicationic ionic liquids encompassing pyridinium hydrazone-phenoxy conjugates as antimicrobial agents targeting diverse high resistant microbial strains, *J. Mol. Liq.*284 (2019) 431-444.
- [75] O.O. Ajani, K.T. Iyaye, O.Y. Audu, S.J. Olorunshola, A.O. Kuye, I.O. Olanrewaju, Microwave assisted synthesis and antimicrobial potential of quinoline-based 4-hydrazide-hydrazone derivatives, *J. Heterocycl. Chem.*55(1)(2017) 302-312.
- [76] R.A. de Cordeiro, C.V.S. de Melo, F.J. de Marques, R. Serpa, de A.J., J. Evangelista, E.P. Caetano, J. Mafezoli, C.F. de Oliveira, M.R. da Silva, J.P.G. de T. Bandeira, J.L.B. Moreira, R.S.N. Brilhante, M.F.G. Rocha, J.J.C. Sidrim, Synthesis and *in-vitro* antifungal activity of isoniazid-derived hydrazones against *Coccidioides posadasii*, *Microb.*98 (2016) 1-5.
- [77] S. Kethireddy, L. Eppakayala, T.C. Maringanti, Synthesis and antibacterial activity of novel 5,6,7,8-tetrahydroimidazo-[1,2-a]-pyrimidine-2-carbohydrazide derivatives, *Chem. Cent. J.* 9(51) (2015) 4-9.
- [78] Z.A. Kaplancikli, L. Yurttas, G. Turan-Zitouni, A. Ozdemir, G. Goger, F. Demirci, U.A.

- Mohsen, Synthesis and antimicrobial activity of new pyrimidine-hydrazones, *Lett. Drug Des. Discov.* 11 (2014) 76-81.
- [79] U.P. Singh, H.R. Bhat, A. Verma, M.K. Kumawat, R. Kaur, S.K. Gupta, R.K. Singh, Phenylhydrazone bearing pyrazole and pyrimidine scaffolds: Design and discovery of novel class of non-nucleoside reverse transcriptase inhibitors (NNRTIs) against HIV-1 and their antibacterial properties, *RSC Adv.* (2013). <https://doi.org/10.1039/b000000x>.
- [80] P. Sridhar, M. Alagumuthu, S. Arumugam, S.R. Reddy, Synthesis of quinoline acetohydrazide-hydrazone derivatives evaluated as DNA gyrase inhibitors and potent antimicrobial agents, *RSC Adv.* 69 (2013) 1-3.
- [81] T. Govindasami, A. Pandey, N. Palanivelu, A. Pandey, Synthesis, characterization and antibacterial activity of biologically important vanillin related hydrazone derivatives, *IJO.* 1 (2011) 71-77.
- [82] V. Asati, N.K. Sahu, A. Rathore, S. Sahu, D.V. Kohli, Synthesis, characterization and antimicrobial evaluation of some 1,3-benzothiazole-2-yl-hydrazone derivatives, *Arab. J. Chem.* 8 (2015) 495-499. <https://doi.org/10.1016/j.arabjc.2011.01.036>.
- [83] O.O. Ajani, C.A. Obafemi, O.C. Nwinyi, D.A. Akinpelu, Microwave assisted synthesis and antimicrobial activity of 2-quinoxalinone-3-hydrazone derivatives, *Bioorg. Med. Chem.* 18(1) (2010) 214-221.
- [84] A. Deep, S. Jain, P.C. Sharma, P. Verma, M. Kumar, C.P. Dora, Design and biological evaluation of biphenyl-4-carboxylic acid hydrazide-hydrazone for antimicrobial activity, *Acta Pol. Pharm.* 67(3) (2010) 182-183.
- [85] S.A. Khan, P. Mullick, S. Pandit, D. Kaushik, Synthesis of hydrazone derivatives of quinoxalinone-prospective antimicrobial and anti-inflammatory agents, *Acta Pol. Pharm.* 66(2) (2009) 169-172.
- [86] A. Naeem, S.L. Badshah, M. Muska, N. Ahmad, K. Khan, The current case of quinolones: synthetic approaches and antibacterial activity, *Molecules.* 21(4) (2016) 268.
- [87] O.I. El-Sabbagh, H.M. Rady, Synthesis of new acridines and hydrazones derived from cyclic  $\beta$ -diketone for cytotoxic and antiviral evaluation, *Eur. J. Med. Chem.* 44(9) (2009) 3680-3686.
- [88] W. Cai, B. Lian, X. Song, T. Hou, G. Deng, H. Li, A cross-sectional study on mental health among health care workers during the outbreak of Corona Virus Disease 2019, *Asian*



journal of psychiatry. 51 (2020) 102111.

- [89] W.B. Chekol, D.Y. Melesse, Operating room team safety and perioperative anesthetic management of patients with suspected or confirmed novel corona virus in resource limited settings: A systematic review, *Trends Anaesth. Crit. Care.* (2020).
- [90] A.R. Rahmani, M. Leili, G. Azarian, A. Poormohammadi, Sampling and detection of corona viruses in air: A mini review, *Sci. Total Environ.* 740(2020) 140207.
- [91] R. Kharb, M.S. Yar, P.C. Sharma, New insights into chemistry and anti-infective potential of triazole scaffold, *Curr.Med.Chem.* 18(21) (2011) 3265-3297.
- [92] P.C. Sharma, G. Kaur, R. Pahwa, A. Sharma, H. Rajak, Quinazolinone analogs as potential therapeutic agents, *Curr.Med.Chem.* 18(31) (2011) 4786-4812.
- [93] Z. Yang, P. Li, X. Gan, Novel pyrazole-hydrazone derivatives containing an isoxazole moiety: Design, synthesis and antiviral activity, *Molecules.* 23(7) (2018) 1798.
- [94] X.D. Ma, S.Q. Yang, S.X. Gu, Q.Q. He, F.E. Chen, E. de Clercq, J. Balzarini, C. Pannecouque, Synthesis and anti- HIV activity of aryl- 2- [(4- cyanophenyl)amino]- 4- pyrimidinone hydrazones as potent non- nucleoside reverse transcriptase inhibitors, *Chem.Med.Chem.* 6(12) (2011) 2225-2232.
- [95] P. Zhan, H. Liu, X. Liu, Y. Wang, C. Pannecouque, M. Witvrouw, E. De Clercq, Synthesis and anti-HIV activity evaluation of novel N'-arylidene-2-[1-(naphthalen-1-yl)-1H-tetrazol-5-ylthio] acetohydrazides, *Med. Chem.Res.* 19(7) (2010) 652-663.
- [96] P. Vicini, M. Incerti, P.L. Colla, R. Loddo, Anti-HIV evaluation of benzo[d]isothiazolehydrazones, *Eur. J. Med. Chem.* 44(4) (2009) 1801-1807.
- [97] H. Dogan, S.D. Dogan, M.G.Gündüz, V.S.Krishna, C.Lherbet, D.Sriram, O. Sahin, E. Sarıpinar, Discovery of hydrazone containing thiadiazoles as mycobacterium tuberculosis growth and enoyl acyl carrier protein reductase (InhA) inhibitors, *Eur. J. Med. Chem.* 188 (2020) 112035. <https://doi.org/10.1016/j.ejmech.2020.112035>.
- [98] Y.B. Sutar, J.K. Mali, V.N. Telvekar, R.S. Rajmani, A. Singh, Transferrin conjugates of antitubercular drug isoniazid: synthesis and *in vitro* efficacy, *Eur. J. Med. Chem.* 183(2019) 111713. <https://doi.org/10.1016/j.ejmech.2019>.
- [99] E.G. Sampiron, G.F. Costacurta, V.P. Baldin, A.L. Almeida, A.L. Ieque, N.C.S. Santos, V.G. Alves-Olher, F. Vandresen, A.C.R. Gimenes, V.L.D. Siqueira, K.R. Caleffi-Ferracioli, R.F. Cardoso, R.B.L. Scodro, Hydrazone, benzohydrazones and isoniazid-acylhydrazones as

- potential antituberculosis agents. *Future Microbiol.* 14(11) (2019) 981-994. <https://doi.org/10.2217/fmb-2019-0040>.
- [100] S.S. Hakkimane, V.P. Shenoy, S.L. Gaonkar, I. Bairy, B.R. Guru, Antimycobacterial susceptibility evaluation of rifampicin and isoniazid benz-hydrazone in biodegradable polymeric nanoparticles against *Mycobacterium Tuberculosis* H37Rv strain. *Int. J. Nanomedicine*. 13 (2018)4303-4318. <https://doi.org/10.2147/IJN.S163925>.
- [101] S.A. Bonnett, D. Dennison, M. Files, A. Bajpai, T.A. Parish, Class of hydrazones are active against nonreplicating *Mycobacterium Tuberculosis*, *PLoS One*. 13(10) (2018)1-11. <https://doi.org/10.1371/journal.pone.0198059>.
- [102] X. Lu, X. Hu, Z. Liu, T. Zhang, R. Wang, B. Wan, S.G. Franzblau, Q. You, Benzylsulfanyl Benzo-heterocycle amides and hydrazones as new agents against drug-susceptible and resistant: *Mycobacterium Tuberculosis*. *Med. Chem. Comm.* 8(6) (2017) 1303-1306. <https://doi.org/10.1039/c7md00146k>.
- [103] S.D. Joshi, D. Kumar, S.R. Dixit, N. Tigadi, U.A. More, C. Lherbet, T.M. Aminabhavi, K.S. Yang, Synthesis, characterization and antitubercular activities of novel pyrrolyl hydrazones and their Cu-complexes; Elsevier Ltd. 121 (2016). <https://doi.org/10.1016/j.ejmech.2016.05.025>.
- [104] G. Cihan-Ustunda, D. Sātana, G. Ozhan, G. Capan, Indole-based hydrazide-hydrazones and 4-thiazolidinones: synthesis and evaluation as antitubercular and anticancer agents, *J. Enzyme Inhib. Med. Chem.* 31(3)(2016) 369-380. <https://doi.org/10.3109/14756366.2015.1024673>.
- [105] H. A. Abdel-Aziz, W. M. Eldehna, , M. Fares, S. T. A. Al-Rashood, K. A. Al-Rashood, M. M. Abdel-Aziz, D.H. Soliman, Synthesis, biological evaluation and 2D-QSAR study of halophenyl bis-hydrazones as antimicrobial and antitubercular agents, *Int. J. Mol. Sci.* 16(4)(2015) 8719-8743. <https://doi.org/10.3390/ijms16048719>.
- [106] P. Dandawate, K. Vemuri, E. M. Khan, M. Sritharan, S. Padhye, Synthesis, characterization and anti-tubercular activity of ferrocenyl hydrazones and their  $\beta$ -cyclodextrin conjugates, *Carbohydr. Polym.* 108(1)(2014) 135-144. <https://doi.org/10.1016/j.carbpol.2014.03.006>.
- [107] A. Jamadar, A.K. Duhme-Klair, K. Vemuri, M. Sritharan, P. Dandawate, S. Padhye, Synthesis, characterisation and antitubercular activities of a series of pyruvate-containing

- aroylhydrazones and their Cu-complexes, *Dalt. Trans.* 41(30) (2012) 9192-9201.  
<https://doi.org/10.1039/c2dt30322a>.
- [108] A.C. Pinheiro, C.R. Kaiser, T.C.M. Nogueira, S.A. Carvalho, E.F. da Silva, L. de Oliveira Feitosaa, M.D.G.M. de Oliveira Henriques, A.L.P. Candea, M.C.S. Lourencoc, M.V.N. de Souza, Synthesis and antitubercular activity of new L-serinyl hydrazone derivatives, *Med. Chem. (Los. Angeles)*. 7(6) (2011) 611-623.
- [109] P. Kumar, B. Narasimhan, P. Yogeewari, D. Sriram, Synthesis and antitubercular activities of substituted benzoic acid N'-(substituted benzylidene/furan-2-yl-methylene)-N-(pyridine-3-carbonyl)-hydrazides, *Eur. J. Med. Chem.* 45 (12) (2010) 6085-6089.
- [110] R. Bairwa, M. Kakwani, N.R. Tawari, J. Lalchandani, M.K. Ray, M.G.R. Rajan, M.S. Degani, Novel molecular hybrids of cinnamic acids and guanylhyazones as potential antitubercular agents, *Bioorg. Med. Chem. Lett.* 20(5) (2010) 1623-1625.  
<https://doi.org/10.1016/j.bmcl.2010.01.031>.
- [111] A.L.A. Ferreira, A.S. Lopes, A.E.P. Filipe, P.F.E. Pedroso, S.M.A. de Pecorelli, C. Caixado, Quinolyl hydrazones for the treatment of tuberculosis and related diseases, U.S. Patent 9,809,550, B2, November 7 (2017).
- [112] A.L.A. Ferreira, A.S. Lopes, A.E.P. Filipe, C. Caixado, P.F.E. Pedroso, S.M.A. de Pecorelli, Pyridyl hydrazones for the treatment of tuberculosis and related diseases, U.S. Patent 9,593,103, B2, March 14 (2017).
- [113] H.N. Jayaram, P. Kusumanchi, M. Grifantini, P. Franchetti, L. Cappellacci, R. Petrelli, Hydrazone derivatives having potent antitumor activity toward multi-drug resistant tumor cells, U.S. Patent 9,133,230, B2, September 15 (2015).
- [114] A.R. Shelar, M. Merlani, M. Shelar, L. Amiranashvili, B. Shelar, Medicinal applications of benzoic acid hydrazones synthesized on the basis of steroidal tigogenin, U.S. Patent 8,623,849, B2, January 7 (2014).
- [115] T.A.M. Hassan, A.A. Kadi, H.A. Abdel-Aziz, N, N'-Hydrazino-bis-isatin derivatives with selective activity against multidrug-resistant cancer cells, U.S. Patent 8,497,296, B2, July 30 (2013).
- [116] A. Nardi, J.K. Christensen, D. Peters, T. Dyhring, N-Acylhydrazone derivatives useful as modulators of nicotinic acetylcholine receptors, U.S. Patent 8,101,649, B2, January 24 (2012).

- [117] T. Sasaki, H. Lai, N.P. Singh, Conjugates of artemisinin-related endoperoxides and hydrazone derivatives for the treatment of cancer, U.S. Patent 7,692,030, B2, April 6 (2010).
- [118] F. Martin, A. Hoang, Novel silane compounds carrying a hydrazone or diazo functional group in order to functionalize solid supports and immobilize biological molecules on these supports, U.S. Patent 2009/0208374, A1, August 20 (2009).
- [119] S. Kobayashi, Y. Yamashita, H. Ishitani, Process for asymmetric intramolecular (3+2) cyclo-addition of hydrazones, U.S. Patent 7,351,831, B2, April 1 (2008).
- [120] K.N. Winzenberg, A.G. Meyer, Q. Yang, A.G. Riches, N-Phenyl-1,1,1-trifluoromethane-sulfonamide hydrazone derivative compounds and their usage in controlling parasites, U.S. Patent 2007/0238700, A1, October 11 (2007).
- [121] K. Kawagoe, K. Motoki, T. Odagiri, N. Suzuki, C. Chen, T. Mimura, Hydrazone derivatives, U.S. Patent 2006/0276433, A1, December 7 (2006).
- [122] J.K. Almstead, N.J. Izzo, D.R. Jones, R.M. Kawamoto, Medicinal uses of hydrazones, U.S. Patent 6,660,737, B2, December 9 (2003).
- [123] G.A. Reichard, N. Shih, D. Wang, Substituted oximes and hydrazones as neurokinin antagonists, U.S. Patent 6,534,502, B2, March 18 (2003).

**Declaration of interests**

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: